arXiv:hep-ph/0510169v1 13 Oct 2005

Form factors of the $K_{\mu 3}$ decay in the spectator approximation of the Standard Model

V.P. Efrosinin, A.N. Khotjantsev
Institute for Nuclear Research RAS
60 October Revolution Pr. 7A, Moscow 117312, Russia

Abstract

We have calculated the parameter $\xi(0) = f_-(0)/f_+(0)$ of the decay $K^+ \to \pi^0\mu^+\nu_\mu$ in the spectator approximation of the Standard Model. It was obtained that $\xi(0) \simeq -0.02$.

The semileptonic decays of kaons afford an interesting opportunity to study the quark currents. The isotopic symmetry of the strong interactions is successfully used for calculations of such processes. This appears to be an illustration of a definite factorization of the strong and weak interactions at low energies. This circumstance allows us to reveal mechanisms of the quark currents outside the framework of phenomenological approach. The semileptonic decay $K^+ \to \pi^0\mu^+\nu_\mu$ ($K_{\mu 3}$), well studied experimentally, offers an interesting opportunity for such study.

We consider the $K_{\mu 3}$ decay (Fig. 1)

$$K^+(p_K) \to \pi^0(p_\pi) + \mu^+(p_\mu) + \nu_\mu(p_\nu). \tag{1}$$

The tree diagram of this decay is shown in (Fig. 1). The general structure of the Lagrangian of this decay is given by

$$L_W = -\frac{G_F}{\sqrt{2}} sin\theta_C J_l^{+\mu} J_{h\mu}, \tag{2}$$

where $J_l^{+\mu}$ is the weak leptonic current. It reads

$$J_l^{+\mu} = \bar{u}(p_\nu)\gamma^\mu(1-\gamma_5)v(p_\mu), \tag{3}$$

and the phenomenological matrix element of the strangeness changing vector current $J_{h\mu}$ can be expanded in the following form

$$\begin{aligned} J_{h\mu} &= \langle\pi^0|\bar{s}\gamma_\mu u|K^+\rangle \\ &= \frac{1}{\sqrt{2}}[f_+(t)(p_K+p_\pi)_\mu + f_-(t)(p_K-p_\pi)_\mu], \end{aligned} \tag{4}$$



Figure 1: The tree diagram of the $K_{\mu3}$ decay.

where $t = (p_K - p_\pi)^2$ is the four-momentum transfer squared of mesons to the leptons.

It is accepted to consider that in this case only vector part of the weak hadronic current works. The matrix element of the axial-vector current is a pseudovector. It is impossible to construct a pseudovector from two momenta. In the case of the exact SU(3) symmetry, $f_+(0) = 1$ and $f_-(0) = 0$. The physical situation corresponds to the disturbed SU(3) symmetry. The form factors $f_\pm(t)$ are represented as

$$f_\pm(t) = f_\pm(0)(1 + \lambda_\pm \frac{t}{m^2_{\pi^+}}), \tag{5}$$

where parameter $\lambda_+ \simeq 0.028$ [1].

Parameter $\xi(0)$ is determined as the ratio of the two vector form factors

$$\xi(0) = f_-(0)/f_+(0). \tag{6}$$

The value of $\xi(0)$ can be obtained by three different methods: by measuring the muon polarization, by studying the Dalitz plot of the $K_{\mu3}$ decay, and also from the $\Gamma(K_{\mu3})/\Gamma(K_{e3})$ ratio [1]. The consistent experimental values of $\xi \sim -0.1$ are obtained from λ_+, λ_0 and λ_+ - λ_0 measurement [2] and from the ratio $\Gamma(K_{\mu3})/\Gamma(K_{e3})$ [3]. The results of these experiments demonstrate a recovery of the SU(3) symmetry because $\xi(0) \to 0$. But the mechanism of such recovery is not yet well understood.

In the Standard Model (SM), the amplitude of the $K_{\mu3}$ decay is given by

$$\begin{aligned} T &= -\frac{G_F}{2} sin\theta_C \bar{u}(p_\nu)\gamma^\mu(1-\gamma_5)v(p_\mu)[f_+(t)(p_K+p_\pi)_\mu + f_-(t)(p_K-p_\pi)_\mu] \\ &= -G_F sin\theta_C f_+ \bar{u}(p_\nu)[\gamma^\mu(1-\gamma_5)(p_K)_\mu - \frac{1}{2}(\xi(t)-1)m_\mu(1+\gamma_5)]v(p_\mu), \end{aligned} \tag{7}$$

where θ_C is the Cabibbo angle, and G_F is the Fermi coupling constant.

The amplitude of $K_{\mu3}$ decay (7) in the K^+ rest frame can be represented as

$$T = -G_F sin\theta_C f_+ \bar{u}(p_\nu)[\gamma^0(1-\gamma_5)m_K - \frac{1}{2}(\xi(t)-1)m_\mu(1+\gamma_5)]v(p_\mu). \tag{8}$$

In the SM the main contribution to amplitude (7) of the $K_{\mu 3}$ decay at the quark level is described by diagram (Fig. 2). In the spectator model, we consider below, the strange quark $\bar{s}$ interacts with the W^+ boson and transforms into $\bar{u}$ quark. The u quark continues independent motion. The subsequent strong interaction in a final state and fragmentation into a pion do not influence an angular distribution of leptons and the muon polarization. We do not include in this approach possible contributions to $K_{\mu 3}$ decay from various extensions of the SM, because these non-SM models are strongly constrained from the experimental search for T-violation in this decay [4].



Figure 2: The diagram of the $K_{\mu 3}$ decay at quark level.

The amplitude of the diagram (Fig. 2) in the spectator model can be written as

$$T^1 = -\frac{G_F}{\sqrt{2}} sin\theta_C \bar{v}(p_s)\gamma_\mu(1-\gamma_5)v(p_u)\bar{u}(p_\nu)\gamma^\mu(1-\gamma_5)v(p_\mu), \tag{9}$$

where p_s and p_u are the momenta of $\bar{s}$ and $\bar{u}$ quark, respectively.

These objects $\bar{s}$ and $\bar{u}$ can be current quarks, but we assume that they are constituent quarks. Such interpretation is proven due to the experimental test of a number of relations between amplitudes and cross sections. These ratios are derived in the framework of the impulse approximation of the additive quark model [5]. In a simple approach [6], there is a prediction for the existence of a soft scalar interaction of constituent quarks in a hadron, and the spin-spin interaction can be large. For example, it results in a considerable difference of masses of π and ρ mesons. We assume that constituent quarks are non-relativistic. Taking into account the short-distance nature of the weak interaction, we can use an impulse approximation to interacting quarks within the framework of the spectator model.

As it was mentioned above, only vector part of the weak hadronic current contributes to the $K_{\mu 3}$ decay. We will further use the Gordon identity for the vector current of the constituent $\bar{s}$ and $\bar{u}$ quarks

$$\bar{v}(p_s)\gamma_\mu v(p_u) = \frac{1}{m_s + m_u}\bar{v}(p_s)[-(p_s+p_u)_\mu + \sigma_{\mu\nu}(p_s-p_u)^\nu]v(p_u), \tag{10}$$

where

$$\sigma_{\mu\nu} = \frac{1}{2}(\gamma_\mu\gamma_\nu - \gamma_\nu\gamma_\mu),$$

m_s is the rest mass of the constituent $\bar{s}$ quark, and m_u is the rest mass of the constituent $\bar{u}$ quark in pion. It follows from Eq. (10) that the tensor contribution should be added. However, the experimental study of the decay $K^{\pm} \rightarrow \pi^0 e^{\pm}\nu$ strongly constraints the contribution of the tensor interaction to a few per cent level at (90% C.L.) [7, 8]. Therefore, we omit tensor coupling in our approach. So, using Eq. (10), we can present amplitude (9) as

$$T^1 = \frac{G_F}{\sqrt{2}} sin\theta_C \frac{1}{m_s + m_u} \bar{v}(p_s)(p_s + p_u)^\mu v(p_u)\bar{u}(p_\nu)\gamma_\mu(1-\gamma_5)v(p_\mu). \tag{11}$$

In the spectator model

$$(p_s)^\mu = (p_u)^\mu + (p_\mu)^\mu + (p_\nu)^\mu, \tag{12}$$

and

$$\begin{aligned} T^1 &= \frac{G_F}{\sqrt{2}} sin\theta_C \frac{1}{m_s + m_u} \bar{v}(p_s)(2p_s - p_\mu - p_\nu)^\mu v(p_u)\bar{u}(p_\nu)\gamma_\mu(1-\gamma_5)v(p_\mu) \\ &= \frac{G_F}{\sqrt{2}} sin\theta_C \frac{\bar{v}(p_s)v(p_u)}{m_s + m_u} \bar{u}(p_\nu)(\gamma_\mu(1-\gamma_5)2(p_s)_\mu + m_\mu(1+\gamma_5))v(p_\mu). \end{aligned} \tag{13}$$

In the rest frame of K^+ we make the assumption that $p_s = (m_s, 0)$. Then, from expression (13)

$$T^1 = \frac{G_F}{\sqrt{2}} sin\theta_C \frac{\bar{v}(p_s)v(p_u)}{m_s + m_u} \bar{u}(p_\nu)(\gamma^0(1-\gamma_5)2m_s + m_\mu(1+\gamma_5))v(p_\mu). \tag{14}$$

The amplitude (13) must be multiplied by the factored contribution of strong interaction of quarks in the final state. Then, it corresponds to the amplitude (7), and it is correct at small t. Thus, the ratio of scalar and vector parts in (8) is equal to the same ratio in the expression (14)

$$\begin{aligned} &\frac{G_F sin\theta_C f_+ \bar{u}(p_\nu)(1/2)(\xi - 1)m_\mu(1+\gamma_5))v(p_\mu)}{-G_F sin\theta_C f_+ \bar{u}(p_\nu)\gamma^0(1-\gamma_5)m_K v(p_\mu)} \\ = &\frac{(G_F/\sqrt{2}) sin\theta_C(\bar{v}(p_s)v(p_u)/(m_s + m_u))\bar{u}(p_\nu)m_\mu(1+\gamma_5))v(p_\mu)}{(G_F/\sqrt{2}) sin\theta_C(\bar{v}(p_s)v(p_u)/(m_s + m_u))\bar{u}(p_\nu)\gamma^0(1-\gamma_5)2m_s v(p_\mu)}. \end{aligned} \tag{15}$$

From (15) we obtain

$$m_s = \frac{m_K}{1 - \xi}. \tag{16}$$

Such spectator approximation will be correct only for a meson with one heavy quark. For example, for the $\pi^+ \rightarrow \pi^0 e^+ \nu_e$ decay (exact $SU(2)$ symmetry, $\xi(0) = 0$) this approximation is not justified. Therefore, expression (16) has no sense for the precise unitary symmetry and can not be extrapolated to the $\pi^+ \rightarrow \pi^0 e^+ \nu_e$ decay. It is important that the constituent mass of the strange quark is used in this formula.

Let us consider a simple non-relativistic model of constituent quarks with a spin-spin interaction for mesons [6]:

$$V_s = \beta_M \frac{\mathbf{S}_1 \mathbf{S}_2}{m_1 m_2}, \tag{17}$$

where V_s is the spin-spin interaction, $\mathbf{S}_i$ is the spin of constituent quark, m_i is the mass of constituent quark, β_M is the constant of the spin-spin interaction,

$$\begin{aligned} m_\pi &= 2m_u + E_0^M - (3/4)\beta_M m_u^{-2}, \\ m_\rho &= 2m_u + E_0^M + (1/4)\beta_M m_u^{-2}, \\ m_K &= m_u + m_s + E_0^M - (3/4)\beta_M m_u^{-1} m_s^{-1}, \\ m_{K^*} &= m_u + m_s + E_0^M + (1/4)\beta_M m_u^{-1} m_s^{-1}. \end{aligned} \tag{18}$$

Here, E_0^M is the scalar interaction, which is supposed to be small. Then, from (18) we have

$$m_u \simeq \frac{m_\pi + 3m_\rho}{8} \simeq 0.305 \text{ GeV}, \tag{19}$$

$$m_u + m_s \simeq \frac{m_K + 3m_{K^*}}{4} \simeq 0.792 \text{ GeV}. \tag{20}$$

Using equations (16,19,20) one can obtain

$$\xi = 1 - \frac{m_K}{m_s} \simeq -0.02. \tag{21}$$

The accuracy of ξ in this approach is about 0.1 and determined by the uncertainty of the constituent s-quark mass of the model [6]. From the expression (16), the mechanism of the recovery of the unitary symmetry relative to the value ξ is seen. The extraction of $\xi(0)$ with small uncertainty is possible in the experiment E246 [9], where the normal polarization of the muon in the $K_{\mu3}$ decay is expected to be measured with high statistics and small systematic errors. The further precise measurement of ξ in the $K_{\mu3}$ decay will allow us to establish the mechanism of quark currents of week interactions.

References

[1] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.

[2] O. P. Yushchenko *et al.*, Phys. Lett. B **581** (2004) 31 [arXiv:hep-ex/0312004].

[3] K. Horie *et al.* Phys. Lett. B **513** (2001) 311 [arXiv:hep-ex/0106006].

[4] M. Abe *et al.*, Phys. Rev. Lett. **83**, 4253 (1999); M. Abe *et al.*, Phys. Rev. Lett. **93** 2004 131601 [arXiv:hep-ex/0408042].

[5] V. M. Shekhter, Sov. J. Nucl. Phys. **33** (1981) 426 [Yad. Fiz. **33** (1981) 817].

[6] V. P. Efrosinin and D. A. Zaikin, Yad. Fiz. **37** (1983) 1532.

[7] O. P. Yushchenko *et al.*, Phys. Lett. B **589** (2004) 111 [arXiv:hep-ex/0404030].

[8] S. Shumizu *et al.* Phys. Lett. B **495** (2000) 33.

[9] M. Abe *et al.*, Nucl. Instrum. Meth. A **506** 2003 60 [arXiv:hep-ex/0302001].